arXiv:1004.5249v1 [math.AT] 29 Apr 2010

The HELP-Lemma And Its Converse In Quillen Model Categories

R.M. Vogt

April 30, 2010

Abstract We show that a map $p : X \to Y$ between fibrant objects in a closed model category is a weak equivalence if and only if it has the right homotopy extension lifting property with respect to all cofibrations. The dual statement holds for maps between cofibrant objects.

The HELP-Lemma states that a homotopy equivalence $p : X \to Y$ of topological spaces has the homotopy extension lifting property , HELP for short, for all closed cofibrations [2, Appendix Thm. 3.5]. The lemma, variants of it and their Eckmann-Hilton duals (e.g. see [1, II.1.11], [4, Thm. 4, Thm. 4*]) have proven to be very useful tools in homotopy theory. Surprisingly it has been overlooked that the homotopy extension lifting property for all closed cofibrations characterises homotopy equivalences:

Proposition: A map of topological spaces $X \to Y$ is a homotopy equivalence if and only if it has the HELP for all closed cofibrations.

In this note we will prove this statement and its Eckmann-Hilton dual in arbitrary closed model categories $\mathcal{M}$ in the sense of Quillen [5], (see also [3]). Applying our result to the category $\mathcal{T}op$ of toplogical spaces with the Strøm model structure [6] we retrieve the Proposition.

Definition: Let $i : A \to B$ and $p : X \to Y$ be maps in $\mathcal{M}$.

(1) We say that p has the right HELP with respect to i, if for each not

necessarily commutative square

$$\begin{tikzcd} A \arrow[r, "f_A"] \arrow[d, "i"'] & X \arrow[d, "p"] \\ B \arrow[r, "g"] & Y \end{tikzcd} \qquad (*)$$

and each right homotopy $h_A : A \to Y^I$ from $p \circ f_A$ to $g \circ i$, where Y^I is a path object $Y \xrightarrow[\sim]{j} Y^I \overset{\pi}{\twoheadrightarrow} Y \times Y$ for Y, there is a map $f : B \to X$ and a right homotopy $h : B \to Y^I$ from $p \circ f$ to g such that $f \circ i = f_A$ and $h \circ i = h_A$.

(2) We say that i has the left HELP with respect to p, if for each not necessarily commutative square

$$\begin{tikzcd} A \arrow[r, "f"] \arrow[d, "i"'] & X \arrow[d, "p"] \\ B \arrow[r, "g_Y"] & Y \end{tikzcd}$$

and each left homotopy $h_Y : Z_A \to Y$ from $g_Y \circ i$ to $p \circ f$, where Z_A is a cylinder object $A \sqcup A \overset{j}{\rightarrowtail} Z_A \xrightarrow[\sim]{\sigma} A$ for A, there is a map $g : B \to X$ and a left homotopy $h : Z_A \to X$ from $g \circ i$ to f such that $p \circ g = g_Y$ and $p \circ h = h_Y$.

Theorem: (1) A map $p : X \to Y$ of fibrant objects is a weak equivalence in $\mathcal{M}$ if and only if it has the right HELP with respect to all cofibrations.

(2) A map $i : A \to B$ of cofibrant objects is a weak equivalence in $\mathcal{M}$ if and only if it has the left HELP with respect to all fibrations.

Beweis: The two statements are dual so we just prove the first one.

Since X and Y are fibrant the projections

$$\begin{tikzcd} X & X \times Y \arrow[l, "p_X"'] \arrow[r, "p_Y"] & Y \end{tikzcd} \qquad \text{and} \qquad \begin{tikzcd} Y & Y \times Y \arrow[l, "p_1"'] \arrow[r, "p_2"] & Y \end{tikzcd}$$

are fibrations.

Suppose that p is a weak equivalence and that we are given a square $(*)$.

Consider the commutative diagram

$$\begin{tikzcd}
X \arrow[rr, "p"] \arrow[dr, "r"] \arrow[dddr, "\mathrm{id}"'] & & Y \arrow[d, "j"', "\sim"] \arrow[ddr, "\mathrm{id}"] & \\
& P \arrow[r, "q_2"] \arrow[d, two heads, "q_1"] \arrow[dr, phantom, "\textit{pullback}"] & Y^I \arrow[d, two heads, "\pi"] & \\
& X \times Y \arrow[r, "p\times\mathrm{id}"] \arrow[d, two heads, "\mathrm{proj}_X"] \arrow[dr, phantom, "\textit{pullback}"] & Y \times Y \arrow[r, two heads, "\mathrm{proj}_2"] \arrow[d, two heads, "\mathrm{proj}_1"] & Y \\
& X \arrow[r, "p"] & Y &
\end{tikzcd}$$

where $r = (\mathrm{id}, j \circ p)$. Since $\mathrm{proj}_1 \circ \pi$ is a weak equivalence, so is $\mathrm{proj}_X \circ q_1$ and hence r. It follows that q_2 and $\mathrm{proj}_2 \circ \pi \circ q_2$ are weak equivalences. Hence $q_3 = \mathrm{proj}_Y \circ q_1 = \mathrm{proj}_2 \circ \pi \circ q_2 : P \twoheadrightarrow Y$ is a weak equivalence. Now consider

$$\begin{tikzcd}
A \arrow[r, "u"] \arrow[d, tail, "i"'] & Q \arrow[r, "\overline{g}"] \arrow[d, two heads, "\sim"', "q_4"] \arrow[dr, phantom, "\textit{pullback}"] & P \arrow[d, two heads, "\sim"', "q_3"] \\
B \arrow[r, "\mathrm{id}"] & B \arrow[r, "g"] & Y
\end{tikzcd}$$

where $u = (i, v)$ and $v : A \to P$ is induced by $(f_A, g \circ i) : A \to X \times Y$ and $h_A : A \to Y^I$. Since q_4 is a trivial fibration there is a section $s : B \to Q$ such that $s \circ i = u$. Define

$$\begin{aligned} &f = \mathrm{proj}_X \circ q_1 \circ \overline{g} \circ s : B \to X \\ &h : q_2 \circ \overline{g} \circ s : B \to Y^I \end{aligned}$$

Conversely, suppose that p has the right HELP. Consider the diagram

$$\begin{tikzcd}
A \arrow[r, "u"] \arrow[d, tail, "i"] \arrow[dr, phantom, "I"] & P \arrow[r, "\mathrm{proj}_X \circ q_1"] \arrow[dr, "q_3"'] & X \arrow[d, "p"] \\
B \arrow[rr, "g"'] & & Y
\end{tikzcd}$$

where square I is supposed to commute. We define

$$f_A = \mathrm{proj}_X \circ q_1 \circ u : A \to X \quad \text{and} \quad h_A = q_2 \circ u : A \to Y^I.$$

Then h_A is a right homotopy from $p \circ f_A$ to $g \circ i$. Hence there exist

$$f : B \to X \quad \text{and} \quad h : B \to Y^I$$

such that h is a right homotopy from $p \circ f$ to g and $f \circ i = f_A$ and $h \circ i = h_A$. Then f and h induce a map

$$k : B \to P$$

such that $k \circ i = u$ and $q_3 \circ k = \mathrm{proj}_2 \circ \pi \circ k = g$.

Hence q_3 has the right lifting property with respect to all cofibration and has to be a trivial fibration. Since $q_2 : P \to Y^I$ is a right homotopy from $p \circ \mathrm{proj}_X \circ q_1$ to q_3 and since a map right homotopic to a weak equivalence is itself a weak equivalence, $p \circ \mathrm{proj}_X \circ q_1$ is a weak equivalence. Since $\mathrm{proj}_X \circ q_1$ is a weak equivalence, so it p. □

References

[1] H.J. Baues, Algebraic homotopy, Cambridge University Press 1989.

[2] J.M. Boardman, R.M. Vogt, Homotopy invariant structures on topological spaces, Lecture Notes in Math. 347, Springer Verlag, Berlin 1973.

[3] W.G. Dwyer, J. Spalinski, Homotopy theories and model categories, Handbook of Algebraic Topology (I.M. James, ed.), Elsevier Science B.V., 1995.

[4] J.P. May, The dual Whitehead theorems, Topological Topics (I.M. James, ed.),London Math. Soc. Lecture Notes Ser. 86, Cambridge University Press 1983.

[5] D.G. Quillen, Rational homotopy theory, Ann. Math. 90 (1969), 205-295.

[6] A. Strøm, The homotopy category is a homotopy category, Arch. Math. 23 (1972), 435-441.